NEMGUARD[®] SC

NEMATICIDE IN CONCENTRATE SUSPECTION (SC)

COMPOSITION: 100 grams of product contain: Garlic extract (purity : 99.9%) g 100.00 DANGER INDICATIONS: H317: May cause an allergic skin reaction. WARNING ADVICE: CAUTION P261: Avoid breathing vapors. P280: Wear protective gloves / protective clothing / Protect the eyelids. P302 + P352: IF SWALLOWED, SKIN: Rinse thoroughly with water and soap. P333 + P313: In case of skin irritation or rash: consult a physician. P362 + P364: Remove all contaminated clothing and wash them before wearing them again. ADDITIONAL INFORMATION: EUH40I: To avoid risks to human health and the environment, follow the instructions for use. SPI: Do not contaminate water with the product or its container. Keep container tightly closed. During normal handling, a strong smell of garlic develops, which can cause effects lacrimation. If this happens, bring the patient to fresh air and, if necessary, wash them Eyes with ac ua. If symptoms persist consult a physician.

CBC (Europe) S.r.l. Via E. Majorana, 2 - 20834 Nova Milanese (MB) - Tel, 0362-365079 OFFICINA DI PRODUZIONE E CONFEZIONAMENTO ECOspray Ltd. - Grange Farm, Cockley Cley Road, Hilborough, Thetford, Norfolk, IP26 5BT, Gran Bretagna OFFICINA DI CONFEZIONAMENTO I.R.C.A. Service S.p.A. - S.S. Cremasca 591, IO- 24040 Fornovo S. Giovanni (BG)

Registration no. 16742 of the Ministry of Health 11 July 2017

Contents: 0.5 - 1 - 5 - 10 - 20 liters

NEMGUARD® is a registered trademark of ECOspray Ltd.

INFORMATION FOR DOCTOR

In case of intoxication call your doctor for the usual first aid measures.

WARNINGS FOR STORAGE

Store the product in its original packaging in a cool, dry place. Protect from frost. TO

Ambient temperature the product has a guaranteed stability of at least 2 years.

CHARACTERISTICS

NEMguard®SC is a nematocide based on garlic extract that acts against free forms (ie Developmental stages in the soil and not within the plant tissues) of Galls nematodes from the genera *Meloidogyne*, and the others genera *Tylenchus*, *Trichodorus*, *Longidorus*, *Pratylenchus*, *Xiphinema*, *Globodera*, *Heterodera*.

APPLICATION AND DOSES OF USE

Apply NEMguard® SC at a dose of 2-4 L / ha to the growing soil using a suitable irrigation system, using water volumes per irrigation cycle between 1,000 and 30,000 L / ha. The best conditions for using the plant protection product are obtained when the soil is moist enough. It is therefore recommended to add the product to irrigation water towards the end of the irrigation cycle.

AUTHORIZED CROPS:

Solanaceae (tomato, peppers, eggplant): 2-4 L / ha. - Field and greenhouse use.

Period of use: from transplantation / sowing to harvesting.

Interval between interventions: 10-14 days.

Maximum number of interventions: 6 per crop cycle.

Cucurbitacee with edible skins (cucumber, zucchini): 2-4 L / ha. - Field and greenhouse use.

Period of use: from transplantation / sowing to harvesting.

Interval between interventions: 10-14 days.

Maximum number of interventions: 6 per crop cycle.

Cucurbitacee with non-edible skins (melon, watermelon, pumpkin): 2-4 L / ha. - Field and greenhouse use.

Period of use: from transplantation / sowing to harvesting.

Interval between interventions: 10-14 days.

Maximum number of interventions: 6 per crop cycle.

COMPATIBILITY '

The product must be used alone.

PRE-HARVEST INTERVAL (PHI) : not necessary.



Authorized label with the decree of the 11th July 2017 no. 16742